			N	AM	E		I.D.
-	(facility)						
Dyskinesia Identification System:				XAN	л тү	PE (ch	scoring
Condensed User Scale (DISCUS) CURRENT PSYCHOTROPICS/ANTI- CHOLINERGIC AND TOTAL MG/DAY mg				4. (5. (6. (Annu Semi D/C - D/C -		nth leading to the control of the co
		•	I -	8. 0			Moderate (abnormal movements occur frequently and are easy to detect)
	m	•		1.1	Vone		(check one) 4 — Severe (abnormal movements occur almost continuously and are easy to detect) NA — Not Assessed (an assessment for an item is not
See	Instructions On Other Side		_	2. F 3. F			able to be made)
·ASS	SESSMENT DISCUS Item and Score (circle one s	core	for e	each	item)	EVALUATION (see other side)
FACE	1. Tics 0	1	2 2	3	4	NA NA	Intensity prerequisite met? : YES NO Greater than 90 days
EYES	3. Blinking 0	1	2	3	4	NA	neuroleptic exposure? : YES NO 3. Other diagnostic conditions? : YES NO (if yes, specify)
ORAL	4. Chewing/Lip Smacking 0 5. Puckering/Sucking/ Thrusting Lower Lip 0	1	2 2 ·	3	4	NA NA	
LINGUAL	7. Tonic Tongue 0 8. Tongue Tremor 0 9. Athetoid/Myokymic/	1		3 3 3	4 4 4	NA NA NA	4. Last exam date : Last total score : Last conclusion : Preparer signature and title for items 1-4 (if different from physician):
HEAD/ NECK/ TRUNK	10. Retrocollis/Torticollis 0 11. Shoulder/Hip Torsion 0	1	2	3	4	NA NA	5. Conclusion (circle one): A. No TD (if intensity pre- D. Withdrawal TD requisite met, list other diagnostic condition or explain in comments) 5. Conclusion (circle one): 6. Persistent TD F. Remitted TD
UPPER	12. Athetoid/Myokymic Finger-Wrist-Arm 0 1 13. Pill Rolling 0 1		2	3 3	4	NA NA	B. Probable TD G. Other (specify in comments) C. Masked TD 6. Comments:
I M	14. Ankle Flexion/ Foot Tapping 0 1 15. Toe Movement 0		2 2	3 3	4	NA NA	
CON	MMENTS/OTHER	- ;	TOT.	RE	I 5 on	ıly)	
				M D	ATE		
RATER	SIGNATURE AND TITLE	_	NEX	(T EX	KAM	DATE	PHYSICIAN SIGNATURE DATE

Simplified Diagnoses for Tardive Dyskinesia

PREREQUISITES. — The 3 prerequisites are as follows. Exceptions may occur.

- A history of at least three months' total cumulative neuroleptic exposure. Include amoxapine in all categories below as well.
- 2. The presence of at least a "moderate" (3) or "severe" (4) movement on any item or at least two "mild" (2) movements on two items located in different body areas.
- 3. Other conditions are not responsible for the abnormal involuntary movements.

DIAGNOSES. — The diagnosis is based upon the current exam and its relation to the last exam. The diagnosis can shift depending upon: (a) whether movements are present or not, (b) whether movements are present for 3 months or more (6 months if on a semi-annual assessment schedule), and (c) whether neuroleptic dosage changes occur and effect movements.

- NO TD. Movements are not present on this exam or movements are present, but some other condition is responsible for them. The last diagnosis must be NO TD, PROBABLE TD, or WITHDRAWAL TD.
- PROBABLE TD. Movements are present on this exam. This is the first time they are present or they
 have never been present for 3 months or more. The last diagnosis must be NO TD or PROBABLE TD.
- PERSISTENT TD. Movements are present on this exam and they have been present for 3 months or
 more with this exam or at some point in the past. The last diagnosis can be any except NO TD.
- MASKED TD. Movements are not present on this exam but this is due to a neuroleptic dosage increase or reinstitution after a prior exam when movements were present. Also use this conclusion if movements are not present due to the addition of a non-neuroleptic medication to treat TD. The last diagnosis must be PROBABLE TD, PERSISTENT TD, WITHDRAWAL TD, or MASKED TD.
- REMITTED TD. Movements are not present on this exam but PERSISTENT TD has been diagnosed and
 no neuroleptic dosage increase or reinstitution has occurred. The last diagnosis must be PERSISTENT TD or
 REMITTED TD. If movements re-emerge, the diagnosis shifts back to PERSISTENT TD.
- WITHDRAWAL TD. Movements are not seen while receiving neuroleptics or at the last dosage level but are seen within 8 weeks following a neuroleptic reduction or discontinuation. The last diagnosis must be NO TD or WITHDRAWAL TD. If movements continue for 3 months or more after the neuroleptic dosage reduction or discontinuation, the diagnosis shifts to PERSISTENT TD. If movements do not continue for 3 months or more after the reduction or discontinuation, the diagnosis shifts to NO TD.

INSTRUCTIONS

- The rater completes the Assessment according to the standardized exam procedure. If the rater also completes Evaluation items 1-4, he/she must also sign the preparer box. The form is given to the physician. Alternatively, the physician may perform the assessment.
- The physician completes the Evaluation section. The physician is responsible for the entire Evaluation section and its accuracy.
- 3. IT IS RECOMMENDED THAT THE PHYSICIAN EX-AMINE ANY INDIVIDUAL WHO MEETS THE 3 PRE-REQUISITES OR WHO HAS MOVEMENTS NOT EX-PLAINED BY OTHER FACTORS. NEUROLOGICAL ASSESSMENTS OR DIFFERENTIAL DIAGNOSTIC TESTS WHICH MAY BE NECESSARY SHOULD BE OBTAINED.
- 4. File form according to policy or procedure.

OTHER CONDITIONS (partial list)

 Age 	•
-------------------------	---

2. Blind

3. Cerebral Palsey

4. Contact Lenses

5. Dentures/No Teeth

6. Down's Syndrome

7. Drug Intoxication (specify)

8. Encephalitis

9. Extrapyramidal Side-Effects (specify)

10. Fahr's Syndrome

11. Heavy Metal Intoxication (specify)

12. Huntington's Chorea

13. Hyperthyroidism

14. Hypoglycemia

15. Hypoparathyroidism

16. Idiopathic Torsion
Dystonia

17. Meige Syndrome

18. Parkinson's Disease

19. Stereotypies

20. Sydenham's Chorea

21. Tourette's Syndrome

22. Wilson's Disease

23. Other (specify)